

OPEN

SUBJECT AREAS:  
COMMUNITY ECOLOGY  
BIOGEOGRAPHY  
MICROBIAL ECOLOGY  
ENVIRONMENTAL  
MICROBIOLOGYReceived  
23 January 2013Accepted  
26 March 2013Published  
10 April 2013Correspondence and  
requests for materials  
should be addressed to  
Y.P. (yxpan@mail.  
iggcas.ac.cn) or K.N.  
(knealson@usc.edu)

# Integrating niche-based process and spatial process in biogeography of magnetotactic bacteria

Wei Lin<sup>1,2</sup>, Yinzhao Wang<sup>1,2</sup>, Yuri Gorby<sup>3</sup>, Kenneth Nealson<sup>3,4</sup> & Yongxin Pan<sup>1,2</sup>

<sup>1</sup>Biogeomagnetism Group, Paleomagnetism and Geochronology Laboratory, Key Laboratory of the Earth's Deep Interior, Institute of Geology and Geophysics, Chinese Academy of Sciences, Beijing 100029, China, <sup>2</sup>France-China Bio-Mineralization and Nano-Structures Laboratory, Institute of Geology and Geophysics, Chinese Academy of Sciences, Beijing 100029, China, <sup>3</sup>Department of Earth Sciences, University of Southern California, Los Angeles, CA 90089, USA, <sup>4</sup>The J. Craig Venter Institute, 10355 Science Center Drive, San Diego, CA 92121, USA.

Microorganisms play key roles in biogeochemical and nutrient cycling in all ecosystems on Earth, yet little is known about the processes controlling their biogeographic distributions. Here we report an investigation of magnetotactic bacteria (MTB) designed to evaluate the roles of niche-based process and spatial process in explaining variation in bacterial communities across large spatial scales. Our results show that both environmental heterogeneity and geographic distance play significant roles in shaping dominant populations of MTB community composition. At the spatial scale in this study, the biogeography of MTB is relatively more influenced by environmental factors than geographic distance, suggesting that local conditions override the effects of dispersal history on structuring MTB community. Of note, we found that the strength of geomagnetic field may influence the biogeography of MTB. We argue that MTB have the potential to serve as a model group to uncover the underlying processes that influence microbial biogeography.

A major challenge in biogeography is to identify the factors that regulate diversity and distribution of organisms on Earth. Although biogeographic patterns for animals and plants are well documented, the diversity and distribution of microorganisms, which play key roles in all ecosystems, are poorly understood. Whether microorganisms represent cosmopolitan or ecologically restricted distribution is a contentious and hotly debated topic<sup>1</sup>. It was previously assumed that microorganisms have a random and cosmopolitan distribution because of their large population numbers, small sizes, short generation times, and high dispersal capabilities<sup>2</sup>. However, with the advent of molecular techniques, a rapidly growing body of evidence suggests that microorganisms may exhibit biogeographic patterns<sup>1,3,4</sup>. Niche-based process and spatial process are two alternative strategies proposed to generate and maintain microbial diversity<sup>1</sup>. The former emphasizes the importance of local environmental conditions and assumes that same environments should support similar microbial communities regardless of geographic distances, the so called “everything is everywhere, but, the environment selects” situation. In contrast, the latter strategy emphasizes the dependence of geographic distances rather than environmental gradients, which is the similar in concept to Hubbell’s neutral theory for macroorganisms that stochastic processes and dispersal limitation affect variation in species composition<sup>5</sup>.

Magnetotactic bacteria (MTB) are diverse microbes united by the ability to form intracellular magnetic crystals of magnetite and/or greigite usually arranged into one or more linear chains<sup>6</sup>. These magnetic inclusions called magnetosomes help these bacteria to sense and swim along the Earth’s magnetic field lines (a behavior known as magnetotaxis)<sup>6</sup>. All known MTB are found within the *Alphaproteobacteria*, *Deltaproteobacteria*, *Gammaproteobacteria*, phylum *Nitrospirae*, or the candidate division OP3<sup>7–10</sup>. MTB are able to accumulate up to 2–3% iron per cell by dry weight, which is several orders of magnitude higher than iron in *Escherichia coli*<sup>11</sup>. Considering their wide distribution in diverse aquatic and sedimentary ecosystems and high intracellular iron content, MTB may have global significance in iron cycling<sup>12,13</sup> as well as bulk magnetization of sediments<sup>14,15</sup>.

Despite of their remarkable magnetic abilities and proposed ecological functions, our understanding of MTB biogeography remains very poor. Although several studies have found that some environmental factors, such as salinity<sup>16,17</sup>, temperature<sup>18</sup>, nitrate<sup>19</sup>, or sulfur compounds<sup>20</sup>, could explain MTB abundance or community differences at local or regional scales, little information is available concerning the biogeography of MTB across large




spatial scales<sup>21</sup>. In this study we have compared the diversity and distribution of MTB communities from different aquatic ecosystems ranging over a large spatial scale (Fig. 1 and Supplementary Table S1 online). The goals of the present study were to (i) describe the large-scale biogeographic pattern of MTB communities, (ii) identify environmental factors that may contribute to the distribution of MTB, and (iii) quantify the relative abundances of niche-based and spatial processes involved with the structuring of MTB communities. These results may provide a starting point for understanding the underlying mechanism(s) leading to the biogeography of these, and perhaps other, microorganisms.

## Results


**Magnetic enrichment of MTB and their phylogenetic diversity.** MTB were discovered from all 16 locations across various ecosystems (Supplementary Table S1 online). Different morphologies of MTB cells were identified, such as cocci, rods, vibrios, and spirilla (Fig. 2). Living MTB cells were concentrated and enriched by taking advantage of their motility and magnetotaxis through the “MTB trap” method<sup>22</sup>. The diversity of enriched bacterial samples was assessed by comparison of 16S rRNA genes. Nearly 700 sequences were retrieved after removing sequences of insufficient quality or potential chimeras. The most highly represented taxa were members of the phylum *Proteobacteria* (> 90%). Other sequences were identified to belong to the phyla *Nitrospirae*, *Bacteroidetes*, TM7, OD1, *Actinobacteria*, *Firmicutes*, or unclassified *Bacteria*. It was noted that some fast-swimming non-magnetotactic bacteria could be collected during the magnetic enrichment<sup>23</sup>. In order to remove

these potential contaminations, sequences most similar to non-magnetotactic organisms were arbitrarily attributed to contaminations and were removed from further analyses. We ended up with a total of 580 sequences, in which bacteria related to the order *Magnetococcales* and the genus *Magnetospirillum* in the *Alphaproteobacteria* were the most dominant groups, representing 72% and 26% of all sequences, respectively. Consistent with previous studies<sup>7</sup>, bacteria related to the order *Magnetococcales* dominated the MTB communities in most sampling locations, while bacteria related to the genus *Magnetospirillum* were the major group in a few locations (e.g., L4, QJC and YYH) (Fig. 3). Sequences belonging to the *Deltaproteobacteria* and the phylum *Nitrospirae* were also detected (Figs. 3 and 4). Sequences in the *Deltaproteobacteria* were identified to affiliate with the orders *Desulfobacterales* and *Desulfovibrionales*, while *Nitrospirae* sequences identified here were related to MTB sequences belonging to groups 1 and 3 as reported previously<sup>9</sup>.

In addition to sequence data generated from 16 locations in this study, we included our previously described data set of MTB communities from 9 locations across northern and southern China<sup>21</sup>, and compared all these MTB communities together (Fig. 1). It is appropriate to combine these two data sets because of similar sampling, enrichment, and experimental approaches performed in these studies. Together, a total of more than 900 MTB sequences from 25 locations were analyzed (Fig. 1). These sequences can be clustered into 170, 114 and 65 operational taxonomic units (OTUs) at 99%, 98% and 95% similarity cutoffs, respectively (Figs. 4 and 5). Rarefaction curves for all samples nearly reached an asymptote, indicating



**Figure 1** | The locations of 16 sampling sites in this study and 9 locations (those with \*) from previous study<sup>21</sup> that are compared here. Distribution pies of distinct OTUs (98% similarity threshold) are shown for each site. The sampling sites are described in more detail in Supplementary Table S1 online. The map was generated using GeoMapApp version 2 (<http://www.geomapp.org/>).



**Figure 2** | Representative transmission electron micrographs of MTB cells retrieved in this study.

that we successfully captured the major extent of MTB diversity (Fig. 5).

**Analysing the biogeographic pattern of the MTB.** To investigate the biogeography of MTB across studied locations, we used two distinct approaches to determine pairwise community similarities between samples: the Sørensen index and the UniFrac index. The Sørensen index is a taxonomy-based approach that assesses community differences at a single level of taxonomic resolution by defining OTUs at an arbitrary sequence similarity level (e.g., 98% in this study)<sup>24</sup>. While, the phylogeny-based UniFrac index measures the overall degree of phylogenetic divergence between sets of communities, which allows us to compare community phylogenies in a more integrated manner than the taxonomy-based approach<sup>25</sup>.


It has been demonstrated that similarities between MTB communities significantly decreased with increasing geographic distance (Fig. 6a and c,  $P < 0.001$ ), reflecting the distance-decay relationship<sup>26</sup>. Thus, geographic distance plays a role in controlling MTB distribution similar to that seen in other microorganisms<sup>27,28</sup>. Changes of MTB community also significantly depend on environmental distance between sites (Fig. 6b and d,  $P < 0.001$ ), indicating that environmental conditions influence MTB species composition as well. In addition, we noted that although a few OTUs are shared by up to 8 locations, nearly 70% of OTUs are endemic, i.e., found at a unique

sample location (Fig. 4). Taken together, these results provide strong evidence that the dominant populations of MTB communities at scales used in this study represent restricted distribution, and both local environment and dispersal history influence their biogeographic pattern. The patterns are similar irrespective of which methods (phylogeny-based UniFrac index or taxonomy-based Sørensen index) are used (Fig. 6).

**Factors influencing MTB biogeography.** Permutation-based multiple regression on distance matrices (MRM) was performed to determine environmental variables that significantly contributed to explain the observed variation in dominant MTB communities (Table 1). When the UniFrac index was considered, the variables that significantly explained MTB patterns were salinity, Eh, sulfate, temperature, and strength of geomagnetic field. When the Sørensen index was used, in addition to the above-mentioned factors, total iron was also found to significantly contribute to variance in MTB communities (Table 1).


**Quantification of the relative role of niche-based and spatial processes.** Regressing community similarity against selected environmental factors and geographic distance is an effective approach to quantify the relative roles of niche-based process and spatial process in control of community composition, and is widely applied in

biogeographic studies of macroorganisms<sup>29</sup>. When this approach was used, it was possible to partition the variation in MTB community distance into four components. As shown in Figure 7, most of the explanatory power was pure environment (25.4% for UniFrac index) or both environmental heterogeneity and geographic distance (MIX) (13.9% for Sørensen index). Pure geographic distance alone explained only a minor portion of the variation in MTB communities (0.7% for UniFrac index and 3.6% for Sørensen index). Not surprisingly, more than half of the total variation (63.6–70.9%) remained unexplained by either measured environmental factors or




**Figure 4** | Heatmap showing the abundance and distribution of operational taxonomic units (OTUs at 98% threshold similarity) for 25 16S rRNA gene clone libraries of MTB communities that are compared in this study. The abundance of each OTU in each library is indicated by different colors. On the left-hand side, a neighbor-joining phylogenetic tree shows the phylogenetic relationship between OTUs.

diversity and/or activities of MTB in the studied locations. There are several possible mechanisms that may account for the influence of geomagnetic field. The strength of geomagnetic field may directly affect the growth, metabolism, swimming behavior or biomineralization of MTB<sup>40,41</sup> and thus plays a role in regulating their community composition. In addition, for life on Earth, the geomagnetic field acts as an important protective barrier against cosmic radiation<sup>42</sup>. Regions of relatively weak geomagnetic field strength are likely to experience an increased influence of cosmic radiation at the Earth's surface, which may affect biological processes of MTB communities. Since our studied samples are all from the Northern Hemisphere, it is necessary in future studies to analyze and compare MTB communities from the Southern Hemisphere, as well as higher latitude regions, and to confirm whether variations of geomagnetic field would affect the global distribution of MTB. In addition, further experimental analyses in lab are also necessary to address the underlying mechanisms of magnetic field effects on MTB activity and/or diversity. To our knowledge, this is the first report of potential effects of the Earth's magnetic field on the biogeography of microorganisms in nature, which may improve our understanding links between variation of the Earth's magnetic field and evolution of life on Earth.



**Figure 5** | Rarefaction curves for sequences at 99%, 98%, and 95% sequence similarity levels, respectively.



**Figure 6 | Correlations of MTB community similarity with geographic and environmental distances.** Both phylogeny-based UniFrac index and taxonomy-based Sørensen index were used as community similarity. Environmental distances are normalized. All correlations are statistically significant ( $P < 0.001$ ).

This study on MTB contributes to the current debate in microbial biogeography about the relative roles of niche-based process and spatial process in structuring microbial communities. Some studies have found that environmental heterogeneity, like pH<sup>43</sup> or salinity<sup>38</sup>, is a primary factor influencing microbial distribution, while others have suggested that the distribution of microbial communities is largely controlled by geographic distance<sup>44–46</sup>. In the present study, MRM-based variation partitioning analyses have quantitatively revealed that pure environmental factors (for UniFrac index) or MIX (for Sørensen index) explain more of the variation in community similarity of MTB than do pure geographic distances (Fig. 7). The fraction of MIX is a consequence of co-variation of environmental and spatial variables in nature, and can be interpreted as a spatially structured environmental condition<sup>47</sup>. Thus a high fraction of MIX suggests that environmental heterogeneity could be of great importance in shaping community composition<sup>48</sup>. It thus appears that the niche-based process has stronger influence on MTB community distribution than the spatial process of dispersal history, which is consistent with several studies that emphasize the importance of local environmental conditions in structuring microbial communities<sup>1,4</sup>.

It is important to recognize that in this study geographic distance plays a minor but significant role that should not be ignored. This result indicates that while microbes are thought to have high dispersal capabilities (e.g., transport by migrating animals or water currents)<sup>3</sup>, spatial process of dispersal limitation and/or historical events still play a role in MTB distribution over the spatial scale considered in this study. A number of studies on macroorganisms have concluded that the relative importance of environment and geographic distance is spatial scale dependent, and a similar conclusion was recently reached for microorganisms as well<sup>28</sup>. Taken

together, our study highlights the importance of integrating both niche-based and spatial processes in investigations of microbial biogeography.

One should be aware that the data presented here are based on magnetic enrichment of MTB cells followed by comparison and classification of 16S rRNA genes. This may introduce some potential biases. For example, some slow motile MTB may not be captured through magnetic enrichment, or those sequences not similar to any known MTB populations that were discounted in this study may be

**Table 1 | Permutation-based multiple regression on distance matrices of MTB community distances, based on the UniFrac index or the Sørensen index, with geographic distance and environmental factors between sampling sites**

Variable	UniFrac index	Sørensen index
	$R^2$	$R^2$
Ln-transformed geographic distance	<b>0.110***</b>	<b>0.175***</b>
Geomagnetic field strength	<b>0.106***</b>	<b>0.128***</b>
Salinity	<b>0.301***</b>	<b>0.123***</b>
Sulfate	<b>0.179***</b>	<b>0.099***</b>
Temperature	<b>0.042**</b>	<b>0.035*</b>
Eh	<b>0.062***</b>	<b>0.026*</b>
Iron	0.015 (n.s.)	<b>0.045*</b>
Nitrate	0.010 (n.s.)	0.011 (n.s.)
Phosphate	0.007 (n.s.)	0.000 (n.s.)
Nitrite	0.024 (n.s.)	0.021 (n.s.)
pH	0.001 (n.s.)	0.002 (n.s.)

Abbreviation: n.s., not significant; \*  $P < 0.05$ ; \*\*  $P \leq 0.01$ ; \*\*\*  $P \leq 0.001$ .

from totally novel MTB strains not yet described. Therefore, further culturing efforts, fluorescence *in situ* hybridization or single-cell analyses will be necessary to better understand the overall diversity of MTB in nature. Nevertheless, in spite of these potential biases, this study represents one of the largest cross-site surveys of MTB biogeography conducted to date, and our results have revealed clear biogeographic patterns of studied MTB communities across different locations, suggesting that 16S rRNA gene analysis of magnetically enriched MTB is still an effective approach to compare the general diversity of dominant MTB communities in nature<sup>21,49</sup>. In addition, identical approach was performed for all samples in this study, which would minimize the potential bias of experimental procedures.

Despite recent progress, describing biogeography of microbial communities and ascertaining the relative importance of different factors that account for these trends remains very difficult. One of the challenges is the enormous diversity of microorganisms in natural environments that is difficult to be fully addressed even using the most advanced sequencing technologies. Hence, our knowledge of the fundamental principles influencing microbial biogeography remains limited. Our results presented here suggest that MTB provide an opportunity to test microbial biogeographic theories. Advantages for choosing MTB for microbial biogeography analysis are: (i) MTB are free-living bacteria that are ubiquitous in diverse sedimentary ecosystems; (ii) living MTB cells can be easily enriched through their active magnetotactic behavior, which should better reflect contemporary diversity because the enriched bacteria are free of dead cells and/or ancient DNA; and (iii) the sequence diversity of MTB communities in nature, compared with the whole bacterial community, is moderate, and therefore is easily handled and can be addressed at a high degree of taxonomic resolution. Therefore, MTB have the potential to serve as a model group to uncover the underlying processes that influence microbial biogeography.

In summary, our results show that major populations of MTB do not randomly distribute at large spatial scales but represent an ecologically restricted distribution. Both environmental heterogeneity and geographic distance contribute to this distribution, indicating that the biogeography of MTB is controlled by a combination of niche-based and spatial processes. Environmental heterogeneity (with or without spatial structure) is found to explain more variation in MTB than pure geographic distance, indicating that contemporary

environmental condition is one of major factors in structuring MTB community composition in nature. The community similarity of MTB significantly correlates with strength of the Earth's magnetic field, which suggests that geomagnetic field may affect the diversity and biogeography of MTB. This study will form the basis of more detailed studies to further define the global biogeography and ecological functions of MTB communities.

## Methods

**Sampling, MTB enrichment and micro copic observation.** Surface sediment samples from sixteen locations were collected across different ecosystems in China and USA (Supplementary Table S1 online). Geographic distances between sampling sites ranged from 0.026 km to 12,240 km. At each sampling site, surface sediments from the top 5–20 cm were collected. The existence of MTB in sediment samples was checked through the “hanging-drop” method<sup>50</sup>. MTB were magnetically enriched using the “MTB trap” method as described previously<sup>19,22</sup>. For TEM observation, 20  $\mu$ l of MTB enrichments were deposited on Formvar-carbon-coated copper grids and were imaged using a JEM-1400 microscope operating at 80 kV (JEOL Corporation, Japan). The rest of the enrichments were frozen at  $-20^{\circ}\text{C}$  prior to molecular analysis.

**Environmental factor analysis.** Several environmental factors of bulk surface sediments were measured. Salinity and pH were measured using a HQ40d salinity meter (HACH, Loveland, Colorado, USA) and a Mettler Toledo Delta 320 pH meter (Mettler-Toledo, Greifensee, Switzerland), respectively. Nitrate, nitrite, sulfate, phosphate, and total iron in pore water (8.6(1 rate 47e)-8.7(as(betalsoate 473(a)-TJzing)-531.2-

agarose gel electrophoresis. Purified PCR products were cloned into the pMD19-T vector (TaKaRa, Dalian, China) and chemically DH5 $\alpha$  competent cells (Tiangen, Beijing, China) by following the manufacture's instructions. Randomly selected clones were sequenced using the 27F primer (Beijing Genomics Institute, Beijing, China).

After removing vector contaminations and low-quality sequences, the rest were screened for chimeras using the Greengenes chimera-check tool (Bellerophon server)<sup>52</sup>. Those sequences which were most similar to non-MTB bacteria but unrelated to known MTB sequences were attributed to potential contaminations by non-





40. Wang, X. K. *et al.* Effects of hypomagnetic field on magnetosome formation of *Magnetospirillum magneticum* AMB-1. *Geomicrobiol. J.* **25**, 296–303 (2008).
41. Pan, Y. *et al.* Reduced efficiency of magnetotaxis in magnetotactic coccoid bacteria in higher than geomagnetic fields. *Biophys. J.* **97**, 986–991 (2009).
42. Steinhilber, F. *et al.* 9,400 years of cosmic radiation and solar activity from ice cores and tree rings. *Proc. Natl. Acad. Sci. U. S. A.* **109**, 5967–5971 (2012).
43. Fierer, N. & Jackson, R. B. The diversity and biogeography of soil bacterial communities. *Proc. Natl. Acad. Sci. U. S. A.* **103**, 626–631 (2006).
44. Whitaker, R. J., Grogan, D. W. & Taylor, J. W. Geographic barriers isolate endemic populations of hyperthermophilic Archaea. *Science* **301**, 976–978 (2003).
45. Reche, I., Pulido-Villena, E., Morales-Baquero, R. & Casamayor, E. O. Does ecosystem size determine aquatic bacterial richness? *Ecology* **86**, 1715–1722 (2005).
46. Green, J. L. *et al.* Spatial scaling of microbial eukaryote diversity. *Nature* **432**, 747–750 (2004).
47. Borcard, D., Legendre, P. & Drapeau, P. Partialling out the spatial component of ecological variation. *Ecology* **73**, 1045–1055 (1992).
48. Smith, T. W. & Lundholm, J. T. Variation partitioning as a tool to distinguish between niche and neutral processes. *Ecography* **33**, 648–655 (2010).
49. Flies, C. B., Peplies, J. & Schöler, D. Combined approach for characterization of uncultivated magnetotactic bacteria from various aquatic environments. *Appl. Environ. Microbiol.* **71**, 2723–2731 (2005).
50. Greenberg, M., Canter, K., Mahler, I. & Tornheim, A. Observation of magnetoreceptive behavior in a multicellular magnetotactic prokaryote in higher than geomagnetic fields. *Biophys. J.* **88**, 1496–1499 (2005).
51. Lin, W., Li, J., Schöler, D., Jogler, C. & Pan, Y. Diversity analysis of magnetotactic bacteria in Lake Miyun, northern China, by restriction fragment length polymorphism. *Syst. Appl. Microbiol.* **32**, 342–350 (2009).
52. DeSantis, T. Z. *et al.* Greengenes, a chimera-checked 16S rRNA gene database and workbench compatible with ARB. *Appl. Environ. Microbiol.* **72**, 5069–5072 (2006).
53. Baker, G. C., Smith, J. J. & Cowan, D. A. Review and re-analysis of domain-specific 16S primers. *J. Microbiol. Methods* **55**, 541–555 (2003).
54. Cole, J. R. *et al.* The Ribosomal Database Project: improved alignments and new tools for rRNA analysis. *Nucleic Acids Res.* **37**, D141–D145 (2009).
55. Tamura, K. *et al.* MEGA5: Molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol. Biol. Evol.* **28**, 2731–2739 (2011).
56. Sinnott, R. W. Virtues of the Haversine. *Sky & Telescope* **68**, 159 (1984).
57. Borgatti, S. P., Everett, M. G. & Freeman, L. C. *Ucinet 6 for Windows: software for social network analysis.* (Analytic Technologies, Harvard, MA, 2002).
58. Rosenberg, M. S. & Anderson, C. D. PASSaGE: pattern analysis, spatial statistics and geographic exegesis. Version 2. *Methods Ecol. Evol.* **2**, 229–232 (2011).

## Acknowledgments

We thank Andrea Cheung, Yuelong Huang, Wayne Harris, Haitao Chen, and Youliang Su for assistance with sample collection; and Greig Paterson for helpful discussions about data analysis. This work was supported by the CAS/SAFEA International Partnership Program for Creative Research Teams (KZCX2-YW-T10), the CAS project, and the NSFC grants 40821091 and 41104041. Wei Lin is grateful to the NASA Planetary Biology Internship for supporting his study in University of Southern California in Los Angeles, CA.

## Author contributions

W.L., Y.P. and K.N. designed research; W.L., Y.W. and Y.G. performed field work; W.L. and Y.W. performed research; W.L. analyzed data; W.L., Y.G., Y.P. and K.N. wrote the paper.

## Additional information

**Supplementary information** accompanies this paper at <http://www.nature.com/scientificreports>

**Competing financial interests:** The authors declare no competing financial interests.

**License:** This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivs 3.0 Unported License. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-nd/3.0/>

**How to cite this article:** Lin, W., Wang, Y., Gorby, Y., Nealson, K. & Pan, Y. Integrating niche-based process and spatial process in biogeography of magnetotactic bacteria. *Sci. Rep.* **3**, 1643; DOI:10.1038/srep01643 (2013).