

# Chinese Loess Plateau vegetation since the Last Glacial Maximum and its implications for vegetation restoration

Wenying Jiang<sup>1\*</sup>, Yufen Cheng<sup>1</sup>, Xiaoxiao Yang<sup>1,2</sup> and Shiling Yang<sup>1</sup>

<sup>1</sup>Key Laboratory of Cenozoic Geology and Environment, Institute of Geology and Geophysics, Chinese Academy of Sciences, Beijing 100029, China; and <sup>2</sup>School of Earth Sciences and Resources, China University of Geosciences, Beijing 100083, China

## Summary

1. China has been investing heavily in afforestation programmes to control soil erosion on the vast Chinese Loess Plateau (CLP). This massive afforestation has led to a considerable increase in forest and a decrease in dust-storm activity in some areas. However, there have also been some negative outcomes, including low tree survival rate, increased soil erosion, exacerbated water shortages and deep soil desiccation. One important explanation for these is the use of inappropriate species because of a lack of knowledge of the natural vegetation in the area, which has been largely destroyed by human activities.

2. Natural vegetation in the most recent warm period (the early–mid-Holocene) can serve as an analogue for the ongoing greening programme, particularly under the global warming scenario. In this study, the natural vegetation of the CLP since the Last Glacial Maximum (LGM) was reconstructed from pollen analyses of six loess sections.

3. Our results show that herbs were dominant both in the cold–dry LGM and the warm–humid early–mid-Holocene. During the LGM, vegetation in the north-eastern CLP mainly consisted of *Artemisia*, *Echinops*-type, *Taraxacum*-type and Chenopodiaceae, and vegetation in the south-eastern CLP was characterized by the same types but with a slightly higher incidence of Poaceae. During the early–mid-Holocene, vegetation was more diverse, with Poaceae, *Artemisia*, *Echinops*-type and Chenopodiaceae dominant in the north-east, and *Pinus*, *Corylus*, Poaceae, *Artemisia* and *Selaginella sinensis* dominant in the south-east.

4. *Synthesis and applications.* The ecological restoration of herbs should be considered a priority, although trees and shrubs have been prioritized previously. To balance environmental conservation and farm-income support objectives, we suggest planting *Corylus*, *Juglans* and *Selaginella sinensis* in the south-eastern Chinese Loess Plateau (CLP) because of their edible value or medicinal properties. Given the considerable prevalence of *Selaginella sinensis* and the Asteraceae family in the pollen records, and their useful medicinal effects, the CLP has great potential to be a centre for Chinese medicinal herb production.

**Key-words:** arid and semi-arid climate, herb, Holocene, Last Glacial Maximum, loess, natural vegetation, pollen records

## Introduction

Loess, a wind-transported accumulation derived from arid inland areas, covers an area of ~440 000 km<sup>2</sup> on the Loess Plateau in north-central China (Liu 1985). The main body of the Chinese Loess Plateau (CLP) is found in the middle

reaches of the Yellow River (Fig. 1) and is characterized by an arid and semi-arid climate. Destruction of vegetation cover on the CLP, as a result of long-term human activities, has resulted in severe soil erosion. About 1–64 billion tonnes of sediment are transported into the Yellow River each year (Liu 1985), raising the riverbed downstream and thereby causing frequent devastating floods. To tackle this issue, China has invested hundreds of billions of Yuan in

\*Correspondence author. E-mail: jiang@mail.iggcas.ac.cn

numerous afforestation campaigns since the end of 1950s (Wang *et al.* 2007; Chen, Shao & Li 2008b).

Afforestation has led to a considerable increase in forest resources and a decrease in dust-storm activity in some areas (Shen *et al.* 2003; Cao 2011). However, it has also caused environmental degradation in the arid and semi-arid regions (Cao 2008; Cao *et al.* 2011a,b), including low survival rate of trees (Guo *et al.* 2001; Wang *et al.* 2007), increased soil erosion (Normile 2007; Wang *et al.* 2010), exacerbated water shortages (Cao, Chen & Yu 2009) and deep soil desiccation (Chen, Shao & Li 2008a; Wang, Liu & Liu 2009).

To improve the tree survival rate, some local govern-

**Table 1.** Study sites together with their geographical and climatic characteristics

Section	Latitude (°N)	Longitude (°E)	Altitude (m a.s.l.)	Mean annual temperature (°C)	Mean annual precipitation (mm)
Jingbian	37.50	108.90	1688	7.8	395
Pingliang	35.54	106.86	1514	8.6	490
Heshui	35.78	108.29	1475	9.1	562
Fujian	36.02	109.30	1226	9.2	600
Jiian	36.11	110.64	990	10.0	580
Xiangfen	35.84	111.45	560	11.5	550

2003), if the current warming rate continues, it will rise to the Holocene Optimum temperature by AD 2100. Therefore, the vegetation present in the early–mid-Holocene is a good reference for potential natural vegetation on the Loess Plateau and will provide valuable insights into the restoration of vegetation for policy makers.

To date, native vegetation on the Loess Plateau, as derived from various records, remains controversial, and assertions have been made that it was predominantly forest (Shi 1981, 1991; An, Feng & Tang 2003; Tang & An 2007; Shang & Li 2010), forest-steppe (Zhu 1983, 1994; Liu *et al.* 1996) or steppe (Sun *et al.* 1997; Xie *et al.* 2002; Li, Zhou & Dodson 2003; Jiang & Ding 2005). At present, the climatic conditions of the Loess Plateau vary significantly between regions, with ~250 mm mean annual rainfall and a mean annual temperature of ~8 °C in the north-east and ~650 mm mean annual rainfall and a mean annual temperature of ~14 °C in the south-east. Therefore, observations at a single site or a few geological samples may not represent the whole Loess Plateau vegetation picture that, overall, and systematic paleovegetation studies are urgently required before a practical vegetation restoration strategy can be implemented. In this study, we present pollen records from six sites across the plateau, with the objective of identifying the natural vegetation types since the Last Glacial Maximum (LGM) and to suggest suitable plant types for vegetation restoration under global warming conditions.

## Materials and methods

### SITES AND STRATIGRAPHY

Loess is a wind-blown, silt-sized material. On the CLP, complete loess sequences consist of over 30 loess (L)–soil (S) couplets, which date back to ~2.8 Ma (Yang & Ding 2010). From the top to the bottom, the loess units are labelled L<sub>1</sub>–L<sub>34</sub>, and the interbedded soils are labelled S<sub>0</sub>–S<sub>33</sub>. Loess beds were deposited during cold-dry glacials, whereas soils developed during warm-humid interglacials (Liu 1985; Kukla 1987). The alternation of loess and soils is evidence of cold-dry and warm-humid climate oscillations during the Quaternary (Kukla 1987; Ding *et al.* 2002; Yang & Ding 2010).

Six loess sections from different topographic and geomorphological units, located at Jingbian, Pingliang, Heshui, Fujian,

Jiian and Xiangfen, were logged (Fig. 1). The Jingbian section, near the margin of the Mu Us desert, is located on ‘Liang’ – a flat ridge covered by thick loess. The other five sites form a west–east transect across the middle of the CLP. The Xiangfen section is situated on the terrace of the Fenhe River, while the Pingliang, Heshui, Fujian and Jiian sections are located on ‘Yuan’ – a high table-land consisting of thick loess. These six sites ensure sufficient spatial coverage and all represent various climatic conditions in the main body of the Loess Plateau. Further details for the sites and their climates are shown in Table 1.

All sections consisted of soil unit S<sub>0</sub> and the upper part of loess unit L<sub>1</sub>. The Holocene soil (S<sub>0</sub>), overlain by modern topsoil, is dark in colour because of its relatively high organic matter content. Loess unit L<sub>1</sub>, yellowish in colour and massive in structure, was deposited during the last glacial period. L<sub>1</sub> can generally be subdivided into five subunits termed L<sub>1-1</sub>, L<sub>1-2</sub>, L<sub>1-3</sub>, L<sub>1-4</sub> and L<sub>1-5</sub> (Yang & Ding 2008). L<sub>1-2</sub> and L<sub>1-4</sub> are well developed soils, and the other subunits are typical loess horizons. Previous studies have shown that L<sub>1-1</sub> was deposited in the MIS 2 (~28–11 ka), which includes the LGM (~26.5–19 ka), S<sub>0</sub> was deposited in the early–mid-Holocene (~11–4 ka), and L<sub>1-2</sub> was deposited in the late MIS 3 (~40–28 ka) (Kukla 1987; Ding *et al.* 2002; Lu, Wang & Wintle 2007).

All the six loess–soil sequences are well exposed in natural outcrops along the walls of gullies. In the field, the loess and soil layers are readily recognizable and laterally traceable for long distances. At each site, a ~1.2-m-wide vertical trench was first excavated along the gully wall, and fresh samples (~400 g each) were then taken at 5–10-cm intervals. To ensure that we used a complete cold-dry climatic cycle for vegetation reconstruction, all the sections were sampled down to loess unit L<sub>1-2</sub>, and a total of 498 samples were collected.

### MEASUREMENTS AND ANALYSES

For stratigraphic correlation and pollen sample selection, grain size was first measured for all samples using a SALD-3001 laser diffraction particle analyser (Shimadzu Corporation, Kyoto, Japan). Ultrasonic pretreatment, with the addition of 20% (NaPO<sub>3</sub>)<sub>6</sub> solution, was used to disperse the samples prior to particle size determination. The analytical procedures used were as detailed by Ding *et al.* (1999). Results showed that soil unit S<sub>0</sub> and the well developed soil L<sub>1-2</sub> are consistently finer grained than loess unit L<sub>1-1</sub> (Fig. 2). The correlation of the lithostratigraphic (Fig. 3) and grain-size curves (Fig. 2) between sections indicates the continuity of the loess deposits.

A total of 97 samples were selected for pollen analysis (Fig. 2). Samples from the coarse-grained unit L<sub>1-1</sub> represent deposition during a cold and dry glacial period, while those from the

S S 0 S  
 L1-1 L1-1 L1-1 L1-1  
 L1-2 L1-2 L1-2 L1-2

fine-grained soil units (S<sub>0</sub> and L<sub>1-2</sub>) represent deposition during relative warm and humid interglacial intervals. Pollen grains were extracted using the heavy-liquid method (Li *et al.* 2006) and identified at ×400 magnification using a Nikon ECLIPSE 50i microscope (Nikon Corporation, Tokyo, Japan). For each sample, 80–400 land pollen grains were counted. Percentages of pollen, spores and algae were calculated relative to the sum of all land pollen counted. Pollen diagrams were drawn using the Psimpoll v2.5 program (Beckers 2005) (Queen's University Belfast, UK).

**Results**

The major pollen assemblages found in the loess and loess-paleosol units in each loess section are detailed below.

**JINGBIAN SECTION**

L<sub>1-2</sub>: Arboreal pollen (AP) percentages are low. *Taraxacum*-type pollen dominates herbaceous pollen (HP) assemblages (Fig. 4a).

L<sub>1-1</sub>: a decrease in *Taraxacum* pollen. An increase in pollen of *Artemisia*, *Echinops*, *Chenopodiaceae* and *Caryophyllaceae*.

S<sub>0</sub>: *Artemisia* pollen predominates (80–90%) with a decrease in *Chenopodiaceae* and a slight increase in *Fabaceae*.



Modern topsoil: an increase in the pollen of *Oleaceae*, *Brassicaceae* and *Eragopyrum*.

**PINGLIANG SECTION**

L<sub>1-1</sub>: AP (including *Quercus*, *Ulmus* and *Nitella*) pollen is dominant. *Taraxacum*-type pollen still dominates herbaceous pollen assemblages.

S<sub>0</sub>: *Artemisia* pollen predominates (80–90%) with a decrease in *Chenopodiaceae* and a slight increase in *Fabaceae*.

Modern topsoil: an increase in the pollen of *Oleaceae*, *Brassicaceae* and *Eragopyrum*.



 modern topsoil
  loess
  paleosol
  weakly developed paleosol

#### HESHUI SECTION

L<sub>1-1</sub>: AP is rare (Fig. 4c). Pollen of *Artemisia*, *Taraxacum*-t pe, *Echinops*-t pe and Chenopodiaceae dominates the spectrum.

S<sub>0</sub>: *Corylus* pollen increases to 17%, making a large contribution to AP. *Artemisia* pollen dominates HP. A decrease in the pollen of *Echinops*-t pe, *Taraxacum*-t pe and Chenopodiaceae.

#### FUXIAN SECTION

L<sub>1-2</sub> and L<sub>1-1</sub>: AP <2% (Fig. 4d). HP predominates, including mainl *Artemisia*, *Echinops*-t pe, *Taraxacum*-t pe, Chenopodiaceae and Poaceae.

S<sub>0</sub>: AP increases to 14.5%, mainl because of an increase in *Corylus* (to 13.8%). *Artemisia* pollen increases to 80%. Pollen of *Echinops*-t pe, *Taraxacum*-t pe and Chenopodiaceae decreases. Ferns and algae are also seen, including Pol podiaceae, *Selaginella sinensis* and *Z* gnema.

Modern topsoil: A decrease in the pollen of *Corylus* and *Artemisia*. An increase in *Echinops*-t pe, *Taraxacum*-t pe, Chenopodiaceae, Poaceae, *Fagopyrum* and Solanaceae pollen, and *Selaginella sinensis* spores.

#### JIXIAN SECTION

L<sub>1-2</sub>: AP is as high as 55%, mainl derived from *Cotoneaster* (Fig. 4e). HP consists mainl of *Artemisia*, *Echinops*-t pe, *Taraxacum*-t pe, Chenopodiaceae and Labiatae.

L<sub>1-1</sub>: *Cotoneaster* pollen disappears. Increase in pollen of *Pinus*, *Corylus* and *Forsythia*. HP content increases because of increases in *Artemisia*, *Echinops*-t pes and Chenopodiaceae.

S<sub>0</sub>: AP increases, mainl because of an increase in *Pinus*. *Artemisia* pollen dominates HP. Poaceae pollen increases to 13%. Pollen of *Echinops*-t pes and Chenopodiaceae decreases. Large increase in both *Selaginella sinensis* and *Z* gnema.

#### XIANGFEN SECTION

L<sub>1-2</sub>: AP is lo (<5%), and includes *Pinus* and Oleaceae (Fig. 4f). Pollen of *Artemisia* and *Echinops*-t pe dominates the spectrum.

L<sub>1-1</sub>: Slight increase in AP thanks to increases in *Pinus*, *Quercus*, Oleaceae and *Forsythia*. *Artemisia* pollen dominates the spectrum. Decrease in *Echinops*-t pe pollen.

S<sub>0</sub>: AP increases in both diversity and abundance. Its content increases to 28%, mainly because of an increase in *Pinus*. In total, twelve AP types may be identified. Increase in *Artemisia* and Poaceae pollen and *Selaginella sinensis* spores to 72%, 20% and 41%, respectively. Fall in *Echinops*-type pollen to 4%.

A total of 54 pollen types may be identified in the six loess sections. It is notable that *Artemisia* pollen dominates all the spectra. A detailed comparison of the pollen assemblages between L<sub>1-1</sub> (from a cold and dry period) and S<sub>0</sub> (from a warm and humid period) is given in Table 2.

### Natural vegetation in the geological past

#### VEGETATION IN THE COLD-DRY PERIOD (L<sub>1-1</sub>)

All sections from the CLP show a similar pollen assemblage for the cold-dry period, characterized mainly by *Artemisia*, together with other significant pollen types including *Taraxacum*-type, *Echinops*-type and Chenopodiaceae. These data indicate the presence of steppe vegetation on the CLP during the LGM.

Spatial differences in pollen assemblages can also be observed. First, the content of Chenopodiaceae generally decreases from 20 to 30% in the north-east (Jingbian and Heshui) (Fig. 4a, c) to 5–12% in the south-east (Fuqian and Xiangfen) (Fig. 4d, f). Second, the Poaceae content increases significantly in a south-easterly direction. In northern China, Chenopodiaceae species predominate in desert shrub, and Poaceae are dominant in steppe

(Editorial Committee of Vegetation Map of China, Chinese Academy of Sciences 2007). We therefore infer that during the LGM, there was a desert steppe in the north-east of the CLP and dry steppe in the south-east.

#### VEGETATION IN THE WARM-HUMID PERIOD (S<sub>0</sub>)

Pollen types were much more diverse during the warm-humid period than during the cold-dry period. There was a greater incidence of AP, and less HP. *Artemisia* was still dominant in pollen assemblages. The incidence of *Taraxacum*-type, *Echinops*-type and Chenopodiaceae pollen decreased, while Poaceae pollen became more prevalent. In addition, the percentages of *Selaginella sinensis* and *Zingnema* increased. In general, steppe vegetation still prevailed on the CLP during the early-mid-Holocene.

The spatial differences in pollen assemblages are distinct. The AP content increases to the south-east, and HP decreases accordingly. In the Jingbian and Pingliang sections in the north-eastern CLP, although the AP consists of about ten pollen types (*Pinus*, *Betula*, *Quercus*, etc.), its incidence is very low (<10%) (Fig. 4a, b). In the middle part of the CLP (Heshui and Fuqian), AP (mainly *Corylus*) has a percentage of ~15% (Fig. 4c, d), but this increases to 17–30% (mainly *Pinus*) in the south-east (Jiqian and Xiangfen) (Fig. 4e, f). Heterophilous ferns and algae, such as *Selaginella sinensis* and *Zingnema*, are rare in the north-east (Jingbian, Pingliang and Heshui) (Fig. 4a–c), but their incidence increases to as high as 10–50% in the south-east (Fuqian, Jiqian and Xiangfen)

**Table 2.** Pollen results from the six loess sections

Site	Dry and cold period (L <sub>1-1</sub> )	Warm and humid period (S <sub>0</sub> )
Jingbian	AP (<20%) mainly includes <i>Pinus</i> and <i>Quercus</i> . HP (>80%) mainly consists of <i>Taraxacum</i> -type, <i>Artemisia</i> and Chenopodiaceae.	Small amounts of AP (mainly <i>Pinus</i> and <i>Quercus</i> ). <i>Artemisia</i> is predominant. Chenopodiaceae decreases and Fabaceae increases slightly.
Pingliang	AP (<5%) includes <i>Pinus</i> , <i>Ephedra</i> , <i>Betula</i> , <i>Corylus</i> , <i>Quercus</i> , <i>Juglans</i> , Oleaceae, <i>Ulmus</i> and <i>Nitraria</i> . HP mainly consists of <i>Artemisia</i> , <i>Echinops</i> -type and <i>Taraxacum</i> -type.	AP is rare. <i>Artemisia</i> still dominates. Both <i>Echinops</i> -type and <i>Taraxacum</i> -type decrease.
Heshui	Few AP types. <i>Artemisia</i> is dominant. Other HPs mainly consist of <i>Taraxacum</i> -type, <i>Echinops</i> -type and Chenopodiaceae.	<i>Corylus</i> content rises to 17% (a large contribution to the AP). <i>Artemisia</i> still dominates. <i>Taraxacum</i> -type, <i>Echinops</i> -type and Chenopodiaceae decrease.
Fuqian	AP (<2%) consists of <i>Pinus</i> , <i>Ephedra</i> , <i>Betula</i> , <i>Corylus</i> , Caprifoliaceae, <i>Nitraria</i> and <i>Ulmus</i> . HP is predominant, mainly including <i>Artemisia</i> , <i>Echinops</i> -type, <i>Taraxacum</i> -type, Chenopodiaceae and Poaceae.	AP increases to 14.5%, mainly due to a rise in <i>Corylus</i> (up to 13.8%). <i>Artemisia</i> content is up to 80%. <i>Echinops</i> -type, <i>Taraxacum</i> -type and Chenopodiaceae decrease. Polypodiaceae, <i>Selaginella sinensis</i> and <i>Zingnema</i> also occur.
Jiqian	AP (<18%) includes <i>Pinus</i> , <i>Corylus</i> and <i>Forsythia</i> . <i>Artemisia</i> is dominant. Other HPs mainly consist of <i>Echinops</i> -type, <i>Taraxacum</i> -type and Chenopodiaceae.	AP increases, mainly due to a rise in <i>Pinus</i> . <i>Artemisia</i> still dominates. Poaceae increases. <i>Echinops</i> -types and Chenopodiaceae decrease. Both <i>Selaginella sinensis</i> and <i>Zingnema</i> increase rapidly.
Xiangfen	AP (<10%) includes <i>Pinus</i> , <i>Quercus</i> , Oleaceae, <i>Forsythia</i> and <i>Sorbaria</i> . HP consists of high percentages of <i>Artemisia</i> (50–60%) and <i>Echinops</i> -type (20–40%), and small amounts of <i>Taraxacum</i> -type, Chenopodiaceae and Poaceae.	Twelve AP types. AP increases to 28%, mainly due to a rise in <i>Pinus</i> . <i>Artemisia</i> , Poaceae and <i>Selaginella sinensis</i> increase to 72%, 20% and 41%, respectively. <i>Echinops</i> -type drops to 4%.

AP, Arboreal pollen; HP, Herbaceous pollen.



(Fig. 4d–f). It is therefore clear that during the warm-humid period, meadow-steppe vegetation dominated the south-eastern CLP, while dry steppe prevailed in the north-east.

## Discussion

Our study sites represent a range of geomorphological units and are characterized by different climatic conditions (Table 1), but they all have very thick loess deposits (60–300 m). Chinese loess is mainly composed of loosely cemented silt (Liu 1985; Yang & Ding 2008), which allows rainwater to infiltrate quickly (Chen, Shao & Li 2008b; Yang *et al.* 2012). In areas with thick loess, therefore, the water in the surface soil is insufficient to maintain forests. Only in areas of thin loess underlain by bedrocks (e.g. deep gullies, incipient floodplains and low river terraces), where the underground water table is relatively high, can trees and shrubs grow under the appropriate conditions.

Our pollen results all show that herbs, rather than trees or shrubs, were dominant on the Loess Plateau in both the cold-dry period and the warm-humid period, which is consistent with pollen data from many other loess sections (Sun *et al.* 1997; Li, Zhou & Dodson 2003; Jiang & Ding 2005; Tang & An 2007; Shang & Li 2010). It should be noted that the pollen records from several sections in river valleys indicate that trees were abundant in the Holocene (An, Feng & Tang 2003; Wu *et al.* 2009; Shang & Li 2010). However, these sections are all located in the riparian zones of first-order tributaries of the Yellow River. In the semi-arid Loess Plateau, these riparian zones cover a very limited area.

Most areas in the Loess Plateau are covered by thick loess (>20 m) (Liu 1985); therefore, our records, together with pollen data from many other sites (Sun *et al.* 1997; Li, Zhou & Dodson 2003; Jiang & Ding 2005; Tang & An 2007; Shang & Li 2010), may well be representative of the entire loess area. Because of this, it appears that priority should be given to planting herbs rather than trees or shrubs in current and future greening programmes.

Most plants selected for past and present afforestation programmes have been trees or shrubs tolerant to arid conditions, including *Robinia pseudoacacia*, *Caragana intermedia*, *Amorpha fruticosa*, *Pinus tabulaeformis*, *Populus davidiana*, *Ulmus pumila* and *Hippophae rhamnoides* (Shi & Yang 2002; Cao 2008). However, these species are rare in the natural vegetation record, except for a few species of *Pinus* and *Ulmus*. Our results show that the native vegetation species on the CLP in the past were mainly from the Poaceae and Asteraceae families, which are well adapted to arid and semi-arid climates and to various topographic and geomorphological units in the loess area. For example, *Agropyron cristatum*, a perennial species from the Poaceae family, is very tolerant of drought, cold and grazing and produces high herbage yields in early spring (a season when feed is in short supply). It also has

a fibrous root system that stabilizes disturbed soil. Members of the Poaceae and Asteraceae families are, therefore, likely to be ideal candidate species for ongoing greening programmes. Choosing the most appropriate species is an urgent task for ecologists.

At present, more than 70 million farmers live on the CLP. An ecological restoration must therefore be combined with income generation for local farmers, to promote sustainable development. According to the early–mid-Holocene pollen records, we suggest *Juglans*, *Corylus* and *Selaginella sinensis* as candidate species for the ongoing greening programmes in areas south-east of Heshui and Fuyuan (Fig. 1). *Juglans* (walnut) and *Corylus* (hazel) seeds are important in the food industry. In a recent study (Ottaggio *et al.* 2008), tannins, including 10-deacetyl baccatin III, baccatin III, paclitaxel C, and 7-epipaclitaxel, were found in *Corylus* shells and leaves. Paclitaxel is a popular and expensive anticancer drug, so *Corylus* can be used as a medicinal source material. *Selaginella sinensis*, a native species of fern, is a traditional Chinese medicine for the treatment of hepatitis, cholecystitis, eczema and burns. In addition, the Asteraceae family, which flourishes on the CLP in both cold and warm conditions, is an important source of medicines (Dharmananda 2012). For example, the most effective antimalarial drug artemisinin is derived from *Artemisia annua* (a member of the Asteraceae family) (Covello 2008). The Loess Plateau, therefore, has tremendous potential to be a source of medicinal plants, and this prospect deserves serious consideration in the greening programme.

In summary, as reviewed by Cao (2011), excessive reliance on afforestation has caused significant negative environmental impacts in northern China. According to our pollen records, priority should be given to herbs (especially the Poaceae and Asteraceae families), rather than trees or shrubs, in current and future greening programmes. Only in the deep valleys, with very thin loess and the riparian zones of large rivers, should some trees and shrubs be considered as candidate species. To combine ecological restoration with economic growth, *Juglans*, *Corylus* and some medicinal plants (e.g. *Selaginella sinensis*) should also serve as useful candidate species for the south-eastern Loess Plateau.

## Acknowledgements

This study was supported by the National Basic Research Program of China (973 Program) (2010CB950204), the Chinese Academy of Sciences (CAS) Strategic Priority Research Program (XDA05120203) and the National Natural Science Foundation of China (41172157 and 40972228). We thank Zuoling Chen, Shaohua Feng, Chenxi Xu and Shujun Zhao for their assistance with field work, and three anonymous reviewers for their critical comments.

## References

- An, C., Feng, Z. & Tang, L. (2003) Evidence of a humid mid-Holocene in the eastern part of Chinese Loess Plateau. *Chinese Science Bulletin*, **48**, 2472–2479.

Bennett, K.D. (2005) Psimpoll and Pscomb programs for plotting and



- Wu, H.N., Ma, Y.Z., Feng, Z.-D., Sun, A.Z., Zhang, C.J., Li, F. & Kuang, J. (2009) A high resolution record of vegetation and environmental variation through the last ~25,000 years in the eastern part of the Chinese Loess Plateau. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **273**, 191–199.
- Xiao, J., Xu, Q., Nakamura, T., Yang, X., Liang, W. & Inouchi, Y. (2004) Holocene vegetation variation in the Daihai Lake region of north-central China: a direct indication of the Asian monsoon climatic history. *Quaternary Science Reviews*, **23**, 1669–1679.
- Xie, S.C., Wang, Z.Y., Wang, H.M., Chen, F.H. & An, C.B. (2002) The occurrence of a grass vegetation over the Chinese Loess Plateau since the last interglacier: the molecular fossil record. *Science in China (D)*, **45**, 53–62.
- Yang, S.L. & Ding, Z.L. (2008) Advance-retreat history of the East-Asian summer monsoon rainfall belt over Northern China during the last two glacial-interglacial cycles. *Earth and Planetary Science Letters*, **274**, 499–510.
- Yang, S.L. & Ding, Z.L. (2010) Drastic climatic shift at ~2.8 Ma as recorded in eolian deposits of China and its implications for redefining the Pliocene-Pleistocene boundary. *Quaternary International*, **219**, 37–44.
- Yang, S.L., Ding, Z.L., Wang, X., Tang, Z.H. & Gu, Z.Y. (2012) Negative  $\delta^{18}\text{O}$ - $\delta^{13}\text{C}$  relationship of pedogenic carbonate from northern China indicates a strong response of  $\text{C}_3/\text{C}_4$  biomass to the seasonality of Asian monsoon precipitation. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **317–318**, 32–40.
- Yu, G., Chen, X., Ni, J., Cheddadi, R., Guiot, J., Han, H., Harrison, S.P., Huang, C., Ke, M., Kong, Z., Li, S., Li, W., Li, P., Liu, G., Liu, J., Liu, Q., Liu, K.-B., Prentice, I.C., Quij, W., Ren, G., Song, C., Sugita, S., Sun, X., Tang, L., Van Campo, E., Xia, Y., Xu, Q., Yan, S., Yang, X., Zhao, J. & Zheng, Z. (2000) Palaeovegetation of China: a pollen data-based synthesis for the mid-Holocene and last glacial maximum. *Journal of Biogeography*, **27**, 635–664.
- Zhu, Z.C. (1983) Range of forest steppe zone in northern Loess Plateau. *Acta Phytocologica Sinica*, **7**, 122–131. (in Chinese).
- Zhu, Z.C. (1994) Basic features of forest steppe in the Loess Plateau. *Scientia Geographica Sinica*, **14**, 152–156. (in Chinese with English abstract).

Received 9 November 2012; accepted 11 January 2013

Handling Editor: Shiqiang Wan